



Digestive Health Center

Medication Information

This applies to patients on anti-platelet, diabetic or iron medications.

Review this medication list for medications you are currently taking.

You may take your usual morning medications (that are not included on the list) on the morning of your procedure. **It is very important to take your chronic pain, high blood pressure, heart and seizure medications that you take daily.**

Medication	Instructions
Miralax and other laxatives	Continue taking as you normally do.
Oral Diabetic Medications	Take half of your usual dose of diabetic medication in the morning on the day prior to the exam. Do not take your evening dose of diabetic medication on the day prior to the exam. Do not take any diabetic medication on the morning of your exam. If you usually do Accu-Chek readings at home, please do so in the morning prior to coming to your appointment on the day of your exam.
Diabetic Medications (including insulin and insulin pumps)	Contact your doctor that manages your diabetes for further instructions.
Lovenox (enoxaparin)	Do not take your Lovenox injection the morning of your exam.
Coumadin (warfarin) (<i>Circle One</i>)	Do Not Hold Hold 3 Days Hold 5 Days
Pradaxa, Xarelto, Eliquis (<i>Circle One</i>)	Do Not Hold Hold 1 Day Hold 2 Days Hold 3 Days
Iron (ferrous sulfate)	Stop taking your iron medication 5 days before your procedure. The day after your procedure, you may resume taking iron supplements.
Aspirin	There is no need to stop taking aspirin before your procedure.
Plavix (clopidogrel), Effient, Brilinta (<i>Circle One</i>)	Do Not Hold Hold 5 Days Hold 7 Days
Phentermine Synergy (<i>Phentermine, Bupropion, and Naltrexone</i>)	Hold 4 days
Exenatide IR (<i>Byetta</i>) Liraglutide (<i>Victoza/Saxenda</i>) Semaglutide (<i>Rybelsus</i>) Dulaglutide (<i>Trulicity</i>) Exenatide ER (<i>Bydureon</i>) Semaglutide (<i>Ozempic/Wegovy</i>) Tirzapatide (<i>Mounjaro/Zepbound</i>)	Clear liquids for 24 hours prior to arrival.